

High-velocity runaway stars from three-body encounters

V.V.Gvaramadze¹, A.Gualandris² and S.Portegies Zwart³

¹Sternberg Astronomical Institute, Moscow State University, Universitetskij Pr. 13, Moscow 119992, Russia
 email: vgvaram@mx.iki.rssi.ru

²Center for Computational Relativity and Gravitation, Rochester Institute of Technology, 85 Lomb Memorial Drive, Rochester NY 14623, USA
 email: alessiag@astro.rit.edu

³ Astronomical Institute ‘Anton Pannekoek’ and Section Computational Science, Amsterdam University, Kruislaan 403, 1098 SJ Amsterdam, the Netherlands
 email: spz@science.uva.nl

Abstract. We performed numerical simulations of dynamical encounters between hard massive binaries and a very massive star (VMS; formed through runaway mergers of ordinary stars in the dense core of a young massive star cluster), in order to explore the hypothesis that this dynamical process could be responsible for the origin of high-velocity ($\geq 200 - 400 \text{ km s}^{-1}$) early or late B-type stars. We estimated the typical velocities produced in encounters between very tight massive binaries and VMSs (of mass of $\geq 200 M_{\odot}$) and found that about 3 – 4 per cent of all encounters produce velocities of $\geq 400 \text{ km s}^{-1}$, while in about 2 per cent of encounters the escapers attain velocities exceeding the Milky Ways’s escape velocity. We therefore argue that the origin of high-velocity ($\geq 200 - 400 \text{ km s}^{-1}$) runaway stars and at least some so-called hypervelocity stars could be associated with dynamical encounters between the tightest massive binaries and VMSs formed in the cores of star clusters. We also simulated dynamical encounters between tight massive binaries and single ordinary $50 - 100 M_{\odot}$ stars. We found that from 1 to $\simeq 4$ per cent of these encounters can produce runaway stars with velocities of $\geq 300 - 400 \text{ km s}^{-1}$ (typical of the bound population of high-velocity halo B-type stars) and occasionally (in less than 1 per cent of encounters) produce hypervelocity ($\geq 700 \text{ km s}^{-1}$) late B-type escapers.

Keywords. Stellar dynamics – methods: N-body simulations – binaries: general – stars: neutron.

1. Introduction

The origin of high-velocity runaway stars can be attributed to two basic processes: (i) disruption of a tight massive binary following the (asymmetric) supernova explosion of one of the binary components (Blaauw 1961) and (ii) dynamical three- or four-body encounters in dense stellar systems (Poveda, Ruiz & Allen 1967). In the first process, the maximum velocity attained by runaway stars depends on the magnitude of the kick imparted to the stellar supernova remnant [either a neutron star (NS) or a black hole (BH)] and for reasonable values of this magnitude, the runaway velocity does not exceed $\sim 200 \text{ km s}^{-1}$ (e.g. Portegies Zwart 2000; cf. Gvaramadze 2009). In the second process, the ejection velocity could be higher. For example, the maximum velocity that a runaway star can attain in binary-binary encounters is equal to the escape velocity from the surface of the most massive star in the binaries and could be as large as $\sim 1400 \text{ km s}^{-1}$ (Leonard 1991).

The recent discovery of the so-called hypervelocity stars (HVSs; e.g. Brown et al. 2005) – the ordinary stars moving with velocities exceeding the Milky Way’s escape

velocity – attracted attention to dynamical processes involving the supermassive BH in the Galactic Centre (e.g. Gualandris, Portegies Zwart & Sipior 2005; Baumgardt, Gualandris & Portegies Zwart 2006). These processes [originally proposed by Hills (1988) and Yu & Tremaine (2003)] can result in ejection velocities of several 1000 km s^{-1} . Similar processes but acting in the cores of young massive star clusters (YMSCs) and involving dynamical encounters with intermediate-mass ($\sim 100 - 1000 M_{\odot}$) BHs (IMBHs) were considered by Gvaramadze, Gualandris & Portegies Zwart (2008) to explain the origin of extremely high-velocity ($\sim 1000 \text{ km s}^{-1}$) NSs (e.g. Chatterjee et al. 2005) and HVSs. Gualandris & Portegies Zwart (2007) proposed that exchange encounters between hard binaries and an IMBH formed in the core of a YMSC in the Large Magellanic Cloud could be responsible for the origin of the HVS HE 0437–5439. A strong support to the possibility that at least some HVSs originate in star clusters rather than in the Galactic Centre comes from the proper motion measurements for the HVS HD 271791, which constrain the birth place of this star to the outer parts of the Galactic disc (Heber et al. 2008).

2. High-velocity runaway stars from three-body encounters

In this paper, we explore the hypothesis (Gvaramadze 2007) that some high-velocity runaway stars could attain their peculiar velocities in the course of strong dynamical encounters between hard massive binaries and a very massive ($\geq 100 - 150 M_{\odot}$) star (VMS), formed in the core of a YMSC through collisions and mergers of ordinary massive stars (e.g. Portegies Zwart & McMillan 2002). In this process, one of the binary components is replaced by the VMS, while the second one is ejected (the so-called exchange encounter), sometimes with a high velocity. Our goal is to check whether or not this process can produce early B-type stars (the progenitors of NSs) with velocities of $\geq 200 - 400 \text{ km s}^{-1}$ (typical of pulsars) and $3 - 4 M_{\odot}$ stars with velocities of $\geq 300 - 400 \text{ km s}^{-1}$ (typical of some late B-type halo stars).

In our study we proceed from the similarity between encounters involving a VMS and those involving an IMBH (the latter process is already known to be able to produce high-velocity runaways; Gvaramadze et al. 2008) and the fact that the radii of VMSs are smaller than the tidal radii of the intruders (the massive binaries), so that the tidal breakup and ejection can occur before the binary components merge with the VMS (Gvaramadze 2007; Gvaramadze et al. 2009). Our study is motivated by the recent evolutionary models of VMSs (Belkus et al. 2007; Yungelson et al. 2008), which suggest that VMSs can lose most of their mass via copious winds and leave behind IMBHs with masses of $\leq 70 M_{\odot}$, which are not large enough to contribute significantly to the production of high-velocity runaway stars (see Gvaramadze et al. 2008). We therefore explore the possibility that a VMS could produce high-velocity escapers (either early or late B-type stars) before it finished its life in a supernova and formed a BH. To check this possibility, we performed numerical simulations of three-body exchange encounters using the `sigma3` package, which is part of the `STARLAB` software environment (McMillan & Hut 1996; Portegies Zwart et al. 2001). For a detailed description of the simulations see Gvaramadze et al. (2009).

First, we focus on exchange encounters producing high-velocity early B-type stars and simulate encounters between a binary consisting of two main-sequence stars with masses $m_1 = 40 M_{\odot}$ and $m_2 = 8 M_{\odot}$ and a massive compact body, either a VMS or an ordinary star of mass of $50 - 100 M_{\odot}$ (the most massive ordinary stars formed in clusters with mass of $\simeq 10^3 - 10^4 M_{\odot}$; e.g. Weidner, Kroupa & Bonnell 2009). In order to maximize the ejection speed, we assume that the binary system is very tight, e.g.




Figure 1. *Left:* Velocity distributions at infinity for escaping ($8 M_{\odot}$) stars in encounters between a binary consisting of a primary star with mass $m_1 = 40 M_{\odot}$ and a secondary star with mass $m_2 = 8 M_{\odot}$, and a single VMS of mass $M_{\text{VMS}} = 50 M_{\odot}$, $100 M_{\odot}$, $500 M_{\odot}$ and $1000 M_{\odot}$ (from left to right). The binary semi-major axis is $a = 0.15 \text{ AU}$. *Right:* The probability of exchange encounters between a $(40, 8) M_{\odot}$ binary (with $a = 0.15 \text{ AU}$) and a single VMS resulting in ejection of the $8 M_{\odot}$ star with a velocity from 200 to 700 km s^{-1} (top to bottom).




Figure 2. *Left:* Velocity distributions at infinity for escaping stars in encounters between a binary consisting of a primary star with mass $m_1 = 10 M_{\odot}$ and a secondary star with mass $m_2 = 4 M_{\odot}$, and a single VMS of mass $M_{\text{VMS}} = 50 M_{\odot}$, $100 M_{\odot}$, $500 M_{\odot}$ and $1000 M_{\odot}$ (from left to right). The binary semi-major axis is $a = 0.06 \text{ AU}$. *Right:* The probability of exchange encounters between a $(10, 4) M_{\odot}$ binary (with $a = 0.06 \text{ AU}$) and a single VMS resulting in ejection of the $4 M_{\odot}$ star with a velocity from 200 to 700 km s^{-1} (top to bottom).

formed by tidal capture. In this case, $a \simeq 3r_1 \simeq 0.15 \text{ AU}$, where r_1 is the radius of the primary star (see Gvaramadze et al. 2009). Fig. 1 (left panel) shows the velocity distribution for $8 M_{\odot}$ escapers. As expected, the more massive VMSs are more likely to eject stars with high velocities. In the right panel of Fig. 1 we show the probability of exchange encounters resulting in ejection of the $8 M_{\odot}$ binary component with a velocity from 200 to 700 km s^{-1} (top to bottom). For $M_{\text{VMS}} \geq 100 M_{\odot}$ about 3 per cent of all encounters produce runaways with peculiar velocities $\geq 400 \text{ km s}^{-1}$. It can be seen that even an ordinary star of mass of $50 M_{\odot}$ can occasionally (in ~ 1 per cent of encounters) produce an escape velocity $\geq 400 \text{ km s}^{-1}$. In order to produce escapers with velocities

typical of HVSS ($V_\infty \geq 700 \text{ km s}^{-1}$), a VMS of several hundred solar masses is required. In the case of a $200 - 300 M_\odot$ VMS, ≥ 2 per cent of all encounters result in an escape velocity of $\geq 700 \text{ km s}^{-1}$. This fraction gradually increases to ≥ 3 per cent for the more massive VMSs.

Next, we consider exchange encounters producing high-velocity late B-type stars and simulated encounters between a very tight binary ($m_1 = 10 M_\odot$, $m_2 = 4 M_\odot$ and $a \simeq 3r_1 \simeq 0.06 \text{ AU}$) and a (very) massive star of mass of $50 - 1000 M_\odot$. The velocity distributions for $4 M_\odot$ escapers are shown in Fig. 2 (left panel) for four different values of the VMS mass: $M_{\text{VMS}} = 50, 100, 500$ and $1000 M_\odot$. The right panel of Fig. 2 shows that for all values of M_{VMS} about 2 per cent of encounters results in peculiar velocities $\geq 300 - 400 \text{ km s}^{-1}$ and that about the same percentage of ejected stars attains velocity $\geq 700 \text{ km s}^{-1}$ if $M_{\text{VMS}} \geq 200 M_\odot$.

We therefore argue that the origin of high-velocity ($\geq 200 - 400 \text{ km s}^{-1}$) early and late B-type runaway stars and at least some HVSS could be associated with dynamical encounters between the tightest massive binaries and VMSs formed in the cores of YMSCs. The future proper motion measurements for HVSS with GAIA will reveal what fraction of these extremely high-velocity stars originated in the Galactic disk.

3. Acknowledgements

We are grateful to L.R.Yungelson for useful discussions. VVG acknowledges the Russian Foundation for Basic Research and the International Astronomical Union for travel grants and the Deutsche Forschungsgemeinschaft for partial financial support. AG is supported by grant NNX07AH15G from NASA. SPZ acknowledges support from the Netherlands Organization for Scientific Research (NWO under grant No. 643.200.503) and the Netherlands Research School for Astronomy (NOVA).

References

- Baumgardt, H., Gualandris, A., Portegies Zwart, S. 2006, *MNRAS*, 372, 174
- Belkus, H., Van Bever, J., Vanbeveren, D. 2007, *ApJ*, 659, 1576
- Blaauw, A. 1961, *Bull. Astron. Inst. Netherlands*, 15, 265
- Brown W.R., Geller M.J., Kenyon S.J., Kurtz M.J. 2005, *ApJ*, 622, L33
- Chatterjee, S. et al. 2005, *ApJ*, 630, L61
- Gualandris, A., Portegies Zwart, S. 2007, *MNRAS*, 376, L29
- Gualandris, A., Portegies Zwart, S., Sipior, M.S. 2005, *MNRAS*, 363, 223
- Gvaramadze, V.V. 2007, *A&A*, 470, L9
- Gvaramadze, V.V. 2009, *MNRAS*, 395, L85
- Gvaramadze, V.V., Gualandris, A., Portegies Zwart, S. 2008, *MNRAS*, 385, 929
- Gvaramadze, V.V., Gualandris, A., Portegies Zwart, S. 2009, *MNRAS*, 396, 570
- Heber, U., Edelmann, H., Napiwotzki, R., Altmann, M., Scholz, R.-D. 2008, *A&A*, 483, L21
- Hills, J.G. 1988, *Nat*, 331, 687
- Leonard, P.J.T. 1991, *AJ*, 101, 562
- McMillan, S.L.W., Hut P. 1996, *ApJ*, 467, 348
- Portegies Zwart, S.F. 2000, *ApJ*, 544, 437
- Portegies Zwart, S.F., McMillan, S.L.W. 2002, *ApJ*, 576, 899
- Portegies Zwart, S.F., McMillan, S.L.W., Hut, P., Makino, J. 2001, *MNRAS*, 321, 199
- Poveda, A., Ruiz, J., Allen, C. 1967, *Bol. Obs. Tonantzintla Tacubaya*, 4, 86
- Weidner, C., Kroupa, P., Bonnell, I. 2009, *MNRAS*, in press (arXiv:0909.1555)
- Yu, Q., Tremaine, S. 2003, *ApJ*, 599, 1129
- Yungelson, L.R., van den Heuvel, E.P.J., Vink, J.S., Portegies Zwart, S.F., de Koter, A. 2008, *A&A*, 477, 223